

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6






Fig.7

【Your Medicine】


Prepared on October 16,2002 , Page 1

Mr. Hichiro Yuyama


【Akineton 1mg】

| | | | | | | | | | | |
|---|---------|-----------|---------|---------------------|---|--|--|--|--|--|
|  | | | | | <p>White,uncoated tablet with cleavage line</p>  | | <p>• This medicine is effective in reducing hand or finger tremors, and the rigidity of body muscles, and improving symptoms such as hinder movement or in which the same position cannot be kept.</p> | | <p>• Be careful when you perform any hazardous activity such as working in dangerous heights or driving a motor vehicle, until it is fully known how you will respond to this medicine.</p> <p>• If you go to an eye doctor for glaucoma, please inform your primary doctor.</p> | |
| After breakfast | | | | | | | | | | |
| On waken-ing | Morning | Noon-time | Evening | Before going to bed | 7-day supply | | | | | |
| | 1 | | | | | | | | | |




【Tenormin 50mg】

| | | | | | | | | | |
|------------------|---------|--------------------------------|---------|---|------------------|---|--|--|--|
| | | White film-coated tablet | |  | | <ul style="list-style-type: none">• This medicine has a hypotensive effect.• This medicine is effective in regulating pulse.• This medicine is effective in preventing angina pectoris attacks. | | <ul style="list-style-type: none">• Be careful when you perform any hazardous activity such as working in dangerous heights or driving a motor vehicle, until it is fully known how you will respond to this medicine. | |
| After breakfast | | | | | | | | | |
| On waken- ing | Morning | Noon-time | Evening | Before going to bed | 28-day supply | | | | |
| | 1 | | | | | | | | |

【PL granules】

| | | | | | | | |
|---|---------|----------------|---------|--|---------------|--|--|
|  | | White granules | | <ul style="list-style-type: none">• This medicine is effective in relieving symptoms such as fever,pain,runny nose,or stuffy nose. | | <ul style="list-style-type: none">• Be careful when you perform any hazardous activity such as driving a motor vehicle,because you may feel sleepy when you take it.• If you have experienced symptoms such as vomiting or skin eruptions after taking medicines such as aspirin, please inform your doctor.• If you go to an eye doctor for glaucoma,please inform your primary doctor. | |
| After breakfast,lunch and supper | | | | | | | |
| On waken-ing | Morning | Noon-time | Evening | Before going to bed | 14-day supply | | |
| | 1 | 1 | 1 | | | | |

【Methycobal tablet 250 μg】

| | | | | | | | | | |
|---|---------|----------------------------|---------|---|---------------|--|--|--|--|
|  | | White, suger-coated tablet | |   | | <ul style="list-style-type: none">• This medicine is effective in promoting the action of enzymes.• This medicine acts on the nervous system to improve symptoms of peripheral nerve disorder such as numbness or pain.• This medicine supplements vitamin deficiency. | | | |
| After breakfast | | | | | | | | | |
| On waken-ing | Morning | Noon-time | Evening | Before going to bed | 30-day supply | | | | |
| | 1 | 1 | 1 | | | | | | |